

The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI), and Osteocalcin in Postmenopausal Women

by Nike Hendrijantini

Submission date: 12-Sep-2019 12:02PM (UTC+0800)

Submission ID: 1171232883

File name: ody_Mass_Index_BMI_,_and_Osteocalcin_in_Postmenopausal_Women.pdf (484.82K)

Word count: 3693

Character count: 20419

The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI) and Osteocalcin in Postmenopausal Women

Nike Hendrijantini^{1*}, Rostiny Alle¹, Rosy Setiawati², Eha Renwi Astuti³ and Manggala Pasca Wardhana⁴

¹Department of Prosthodontics, Faculty of Dental Medicine, Airlangga University, Indonesia

²Department of Radiology, Faculty of Medicine, Airlangga University, Indonesia

³Department of Radiology, Faculty of Dental Medicine, Airlangga University, Indonesia

⁴Department of Obstetrics and Gynecology, Faculty of Medicine, Airlangga University, Indonesia

*Corresponding author: Nike Hendrijantini, Department of Prosthodontics, Faculty of Dentistry, Airlangga University, Surabaya, Indonesia, Tel: 62-81-134-1610; E-mail: hendrijantininike@yahoo.com

Received date: June 15, 2016; Accepted date: July 25, 2016; Published date: August 01, 2016

Copyright: © 2016 Hendrijantini N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Osteoporosis is related to the decrease of bone density, osteoporosis requires attention from dentists because it can also occurs in the jaw bone. Osteoporosis is calculated with a quantitative assessment of bone density, which is called Bone Mineral Density (BMD). The best imaging modalities to assess bone density are the Dual Energy examination methods X-ray Absorptiometry (DEXA). In addition to DEXA, osteoporosis examination can be performed with measurement of Body Mass Index (BMI), which is a ratio of weight and height. Process of bone formation by osteoblasts can be examined by bone marker such as Osteocalcin, as a parameter (alone or in combination with BMD) to determine metabolic bone disorders during bone formation and bone remodeling (bone turnover).

Aim: To analyze the correlation among BMD, BMI and Osteocalcin in Postmenopausal women, to investigate the possibility of using osteocalcin examination to predict the mandibular bone osteoporosis.

Method: Fifty four women above 51 years old who are at least 1 year postmenopausal, underwent BMD (using DEXA examinations), BMI, and Osteocalcin. The results of each examination were counted, and the correlations among each examinations were evaluated using Spearman's correlation test.

Result: Means of BMD, BMI, and Osteocalcin in postmenopausal women were 1.606, 25.189, and 30.566 respectively. BMD were significantly correlated with BMI (Spearman's rank correlation coefficient $r=0.414$, $p<0.05$). While, BMD were significantly correlated with Osteocalcin ($r=-0.343$, $p<0.05$). Moreover, BMI were significantly correlated with Osteocalcin ($r=-0.274$, $p<0.05$).

Conclusion: There were significant correlations among BMD, BMI, and Osteocalcin results in postmenopausal women. It is concluded that each examination of BMD, BMI, and Osteocalcin can be used to identify the risk of osteoporosis in postmenopausal women. Therefore simple examination of osteocalcin can be used to predict mandibular bone loss.

Keywords: BMD; BMI; Osteocalcin; Osteoporosis; Postmenopausal women

Introduction

Osteoporosis is a disease which requires more attention from the dentists, osteoporosis is related to the decrease of bone density, and it can also occurs in the jaw bone. The osteoporotic jaw bone will experience the decrease of jaw bone density, the increase alveolar jaw bone resorption, and also the increase risk of periodontal disease and tooth loss. It is crucial for dentists and especially prosthodontist to diagnose jaw bone density, especially in postmenopausal women before prosthodontic treatment. There are numerous failure in prosthodontic treatment due to osteoporosis. Osteoporosis will cause mandible resorption, which in turn cause the failure of osteointegration between the implant and the jaw bone [1].

Problem of osteoporosis in postmenopausal women is associated with the decrease of estrogen level. This condition will accelerate bone fragility [2]. World Health Organisation (WHO) defines osteoporosis as a disease that is characterized by low bone mass and microarchitectural destruction of bone tissue, leading to bone fragility and increased risk of fracture. Diagnosis of osteoporosis according to WHO is obtained from any of the three regions (total hip, femoral neck and lumbar spine). Study Group Meeting of WHO assessed fracture risks and applied it to screen postmenopausal osteoporosis, and recorded that more than 75 million people in the United States, Europe and Japan were suffering from osteoporosis on 2006. Osteoporosis caused more than 8.9 million fractures per year that occurred in the United States and Europe [3]. While, research of Health Department showed that two out of five people in Indonesia have the risk of osteoporosis [4].

Diagnosis of osteoporosis is calculated with a quantitative assessment of bone density, which is called Bone Mineral Density (BMD) [5]. The best imaging modalities to assess bone density are the Dual Energy examination methods x-ray absorptiometry (DEXA) and Quantitative Computerized Tomography (CT). CT is more sensitive but with the added risk of more radiation exposure, several experts prefer DEXA examination due to the less radiation exposure. BMD in postmenopausal women is determined by peak bone mass and the amount of bone lost since menopause [6].

In addition to DEXA, osteoporosis examination can be done with Body Mass Index (BMI). BMI is a ratio of weight and height with units (kg/m^2) [7]. Regarding to BMD and BMI, there are four important factors to assess the quality of bone, these are bone density; bone turnover (remodeling); bone size; and bone geometry (microarchitecture) [6].

Osteocalcin is known as bone Gla protein, a marker of bone formation, vitamin K and vitamin D-dependent protein produced by osteoblasts and most widely studied of the non-collagen protein in bone [8]. Osteocalcin plays an important role in the process of mineralization and calcium ion homeostasis process. Osteocalcin is synthesized during the bone formation and it exhibits a compact, calcium dependent, alpha helical formation, in which the Gama Carboxylglutamic Acid (GLA) residues bind and promote absorption to hydroxyapatite in the bone matrix, thus leading to bone mineralization. Calcium and phosphorus deficient osteoporotic woman may have a decrease rate of bone mineralisation due to a reduction in hydroxyapatite crystal formation [9].

The examination of osteocalcin is a good parameter for determining bone metabolism disorders during bone formation and bone turnover. This marker is often used as an early biomarker of bone-forming therapy and to assess the effectiveness of treatment outcome [6].

Relationship between BMD and BMI is still not clearly defined, some authors found out a positive relationship, while others showed that BMI is a risk factor for osteoporosis. There are some researches showed positive correlation between systemic osteoporosis and the decrease of mandibular bone density [10].

There is increasing evidence that a combination of a biochemical marker and BMD may be a better predictor of fracture than BMD alone, yet biochemical marker such as osteocalcin may also predict fracture risk independently of BMD. High bone turnover can disrupt the trabecular architecture by increasing the incidence of trabecular perforation and buckling, thus reducing bone strength, without necessarily appreciably affecting BMD [9].

In this study we analyzed the correlation among BMD (calculated from lumbar spine), BMI and Osteocalcin in postmenopausal women. While BMD is still the gold standard for osteoporosis diagnosis, the device is unfortunately limited in our country, osteocalcin examination in the other hand is easier, cheaper and more simple. Researcher wants to investigate the possibility of using osteocalcin examination to predict the mandibular bone osteoporosis.

Materials and Methods

Study design

This is a cross sectional observation study, conducted at Pramita and Prodia Laboratory. This study is using fifty four postmenopausal women who received prosthodontic treatment from January until March 2014 in prosthodontic clinic Airlangga University as subjects, and all of them were given informed consent to be examined. Characteristics of subjects were at least 1 year postmenopausal women, above 51 years old, no history of systemic diseases (Hyperparathyroidism, Paget's Disease, Osteomalacia, Renal Osteodystrophy, Osteogenesis Imperfecta), no history of cancer or kidney diseases, no history of hormonal therapy and smoking.

BMD measurement

BMD is the main criteria used to diagnose and monitor osteoporosis. DEXA is the current gold standard examination of osteoporosis in male and female, which is used to examine the bone mass of total hip, femoral neck, lumbar spine, or whole body. The purpose of bone mass measurement are to determine diagnosis of osteoporosis, to predict the occurrence of fractures, to assess changes in bone density after treatment [11]. In this study, position of patients were in lumbar spine with center of laser beam in lumbar. Lumbar spine is measured in this research, due to the faster bone loss nature compared with the hip in post-menopausal women [12]. Measurement of bone density usually expressed in T-scores, where the standard deviations number of bone density varies from the average bone density in normal subjects with the same sex. WHO define osteoporosis as a condition in which bone mineral density is below -2.5 SD, osteopenia in which bone mineral density between -1 to -2.5 SD, while normal in which bone mineral density is above -1 SD [13].

BMI measurement

BMI is a person's weight in kilograms divided by the square of height in meters. Height was measured using a 0.1 microtoise cm precision. Body weight was measured using a digital weight scales Seca brands with 0.1 kg precision. BMI figure obtained by using the weight formula (kg/m^2).

Osteocalcin measurement

The examination of osteocalcin is a good parameter to determine bone metabolism disorders during bone formation and bone turnover. Osteocalcin level were examined using enzyme-linked immunosorbent assay (ELISA). Normal values were: 10.1 ± 9.4 ng/ml [6]. *In vivo* quantitative measurement of N-MID osteocalcin levels were performed in serum or plasma. Examination of the N-Mid Osteocalcin with postmenopausal scale 15-46 ng/ml. Stability samples 2-80°C (immediately after sampling), 200°C (4 hours after sampling, 2 hours if the sample correction carried out in room temperature), -800°C (more than 1 month). This examination is used as a therapeutic control antiresorp. Principles of inspection: The principle of this test is a sandwich ELISA.

In the first incubation, the sample will react with the monoclonal antibody specific N-MID osteocalcin and biotin-labeled monoclonal antibodies specific N-MID osteocalcin labeled ruthenium complex to form a sandwich. Furthermore, the addition of streptavidin-coated

microparticles that cause complex bound to the solid phase via interaction of biotin and streptavidin. The reaction mixture is put into the measuring cell where the microparticles were captured with a magnet on the electrode surface, the compounds are not bound disposed pro cell. Voltage provision to the electrodes induces chemiluminescent emission. Results are determined via a calibration curve made by means of a two-point calibration curves and masks available on the reagent barcode.

Ethical Clearance

This study has recieved ethical clearence from the Comission on Ethics Health Research (KKPEK).

Statistical Analysis

Data analysis was performed using SPSS software (version 21.0; SPSS, Inc., Chicago, IL). Subject characteristics are presented as mean \pm SD unless otherwise stated. Agreement among BMD and BMI, and osteocalcin determined using Spearman's rank correlation coefficient. Significant correlation were found if $p < 0.05$.

Results

Characteristics of participants: There were fifty four postmenopausal women, with average age of 63, 78 years (range 51-83 years old), with 83.33% subjects were in the range of 51-65 years old and 16.67% subjects were in range above 66 years old.

Difference mean of BMD, BMI, and osteocalcin are listed in Table 1. Mean of BMD, BMI, and osteocalcin were 1.606 ± 1.213 , 25.189 ± 4.076 , 30.566 ± 9.144 . Based on BMD value, 27.8% of subjects were osteoporosis, 44.4% of subjects were osteopenia, and 27.8% were normal.

Agreement among BMD, BMI, and osteocalcin with non-parametric method are listed in Table 2, with spearman rank correlation. Assessments of BMD and BMI were significantly related ($p < 0.05$, $r = 0.414$, positive relation). BMD and osteocalcin were significantly related ($p < 0.05$, $r = -0.343$, negative relation), as well as BMI and osteocalcin were also significantly related ($p < 0.05$, $r = -0.274$, negative relation). BMD and BMI had a stronger relation compared with BMI and osteocalcin.

Correlation types are listed in Figure 1. Correlation between BMD and BMI showed linear positive, because dots in graph above were placed in positive axis. Higher BMD linear also has higher BMI. Correlation between BMD and osteocalcin were negative because dots in graph strive were placed in negative axis. Higher BMD has lower Osteocalcin, as well as correlation between BMI and osteocalcin.

Category	Mean	Standard Deviation
BMD	1.606	1.213
BMI	25.189	4.076
Osteocalcin	30.566	9.144

Table 1: Difference Means of BMD, BMI, and osteocalcin with standard deviation.

Correlations					
			BMD	BMI	Osteocalcin
Spearman's rho	BMD	Correlation Coefficient	1	.414**	-.343*
		Significance (2-tailed)	.	0.002	0.011
		N	54	54	54
	BMI	Correlation Coefficient	.414**	1	-.274*
		Significance (2-tailed)	0.002	.	0.045
		N	54	54	54
	Osteocalcin	Correlation Coefficient	-.343*	-.274*	1
		Significance (2-tailed)	0.011	0.045	.
		N	54	54	54
**. Corr. is significant at .01 level 2-tail...					
*. Corr. is significant at .05 level 2-tail...					

Table 2: Correlations among BMD, BMI, and osteocalcin with Spearman correlation statistical test.




Figure 1: Scatterplot correlations from BMD and BMI, BMD and osteocalcin, BMI and osteocalcin.

Discussion

In this study, bone mineral density was evaluated using a DEXA of the spine. Density of the spine is most responsive to drugs intervention and monitoring for patient. DEXA is a technology to determine the relationship between bone density and fracture risk, and has been validated in a longitudinal cohort study. DEXA obtained some information about bone density in a given area in grams/cm², comparison between mean level of bone density compared to the mean level of bone density with the same ethnic adults called T score in percent (%) [11].

This study showed that BMD and BMI had significant correlation (positive correlation) showed in Table 2. This result was consistent with those reported by Hsu et al. [14] and Ahmad et al. [15] which reported a significant positive relationship among weight, BMI and BMD. BMI is a good indicator and often used for measuring BMD [16], this is similar with the research of Iqbal which suggested that BMI is a good indicator for referring patients to do DEXA scanning on Bone Mineral Density (BMD) [17]. The results of the bivariate analysis of Herkulina study showed positive and significant correlation between Body Mass Index (BMI) and Bone Mineral Density (BMD) [18].

The study of Kim et al. also stated that low body weight is more likely associated with osteoporosis [19]. All of height, weight and BMI positively correlated with the dependent variable BMD T-score. As well as research of Hubungan [20] showed that low BMI and decreased bone strength associated with reduced bone mass in all parts of the body. Study of Mendez et al. showed relationship of BMI and BMD was significantly positive at the lumbar spine, pelvis and femoral neck [21].

Correlation between BMD and osteocalcin had significant relationship (negative correlation) showed in Table 1, this is well according to the theory that as a specific marker of osteoblast function, level of osteocalcin have been shown to correlate with bone formation rates, osteoporotic post-menopausal women will have an increase of serum OC concentration. The same statement was made by Kalaiselvi et al. [22] which reported a negative correlation found between the osteocalcin level and the BMD in post-menopausal women (Pearson's correlation $p < 0.05$ and $r = -0.595$). That study in 60 post-menopausal women at Sree Balaji Medical College and Hospital and Bharat Scan Limited at Chennai, India, as well as study of Susanto [9] reported a significant inverse correlation between serum osteocalcin and femoral neck BMD ($p < 0.05$, and $r = -0.29$).

BMI and osteocalcin have significant correlation (negative correlation) showed in Table 2. The same statement is made by Kalaiselvi et al. [22] which reported that a significant correlation was found between osteocalcin level and BMD in post-menopausal women (Pearson's correlation $p < 0.05$, and $r = 0.339$).

Despite BMI usage as a measurement to predict whether a person is overweight or not, it has some drawbacks and may not be an accurate tool for assessing weight status and height. This study is similar with that of body weight or BMI indicates the value inversely with markers of bone turnover, where people with low body weight has higher bone turnover, while overweight and obese have lower serum osteocalcin compared to normal weight. Effect of obesity on the risk of fracture depends on the definition, if defined by BMI or body weight, obesity may protect against bone mineral loss or vertebral fractures. However, if obesity is based on the percentage of body fat, obesity may be a risk factor for osteoporosis [9].

Osteoporosis is not only occurred in lumbar, femur and radius bone, but also on jaw bone, researches have concluded that density of mandible bone related with density of other bones significantly [1]. Based on this research, there was a correlation among BMD, BMI and osteocalcin. Those examinations therefore can be used to predict osteoporosis of the mandible bone.

Conclusion

There were significant correlations among BMD, BMI, and Osteocalcin results in postmenopausal women. BMD and BMI had

positive correlation, while BMD and osteocalcin had negative correlation, and BMI and osteocalcin had also negative correlation. It is concluded that examination of BMD, BMI, and Osteocalcin can be used to identify the risk of osteoporosis in postmenopausal women. Therefore simple examination of osteocalcin can be used to predict bone loss in mandibular bone.

Acknowledgement

To all of respondents whom involved in this study, and also Prodia laboratorium for examining osteocalcin.

References

1. Lindawati SK (2012) Dampak Osteoporosis Tulang Rahang Terhadap Perawatan Prostodontik dan Kualitas Hidup Lansia. *Pidato Pengukuhan Guru Besar FKG UI*, pp. 3-15.
2. Abas BJ dan, Sri P (2007) Risiko Osteoporosis di Indonesia. *Puslitbang Gizi dan Makanan Depkes RI Bogor* 30: 1-11.
3. World Health Organization (2004) 'WHO Scientific Group on The Assessment of Osteoporosis at Primary Health Care Level'. Summary Meeting report Brussel, Belgium, pp.2
4. Ichramsiah AR, dan Bambang S (2010) Penyakit Osteoporosis.
5. Compston J, Cooper A, Cooper C, Francis R, Kanis JA, et al. (2013) Guideline for the diagnosis and management of osteoporosis in postmenopausal women and men from the age of 50 years in the UK. *Maturitas* 62: 105-8.
6. Adi P (2007) Peran Pertanda Tulang Dalam Serum Pada Tata Laksana Osteoporosis. *Universa Medicina* 26: 152-9.
7. Centers For Disease Control (2009) Body Mass Index: 'Considerations for Practitioners. Department of Health and Human Services, USA. pp. 1.
8. Lee AJ, Hodges S, Eatell R (2000) Measurement of Osteocalcin. *Association of Clinical Biochemist* 37: 432-446.
9. Susanto LTM (2011) Serum Osteocalcin and bone mineral density in postmenopausal women. *Universa Medicina* 30: 155-161.
10. Ishii K, Taguchi A, Nakamoto T, Ohtsuka M, Sutthiprapaporn P, et al. (2007) Diagnostic efficacy of alveolar bone loss of the mandible for identifying postmenopausal women with femoral osteoporosis. *Dentomaxillofac Radiol* 36: 28-33.
11. Kawiya IKS (2009) Osteoporosis, Patogenesis Diagnosis dan Penanganan Terkini. *J Penyakit Dalam* 10: 158-162.
12. Bonnick SL, Shulman L (2006) 'Monitoring Osteoporosis Therapy: Bone Mineral Density, Bone Turnover Markers, or Both?'. *The American Journal of Medicine* 119: 255-315.
13. Margo U, Wulandari M, Zilfa KP (2010) Faktor-Faktor Yang Berhubungan Dengan Kepadatan Tulang Pada Wanita Postmenopause. *Jurnal Unimus* 6: 1-2;
14. Hsu YH, Venners SA, Terwedow HA, Feng Y, Niu T, et al. (2006) Relation of body composition, fat mass, and serum lipids to osteoporotic fractures and bone mineral density in Chinese men and women. *Am J Clin Nutr* 83: 146-54.
15. Ahmad EA, Abbas M, Amir R, Mohammad S (2012) 'the Relationship between Body Weight, Body Mass Index (BMI) and Bone Mineral Density (BMD) of the Lumbar Spine and Femoral neck in Professional Cyclist of Iran and Tour de France'. *Iranian Journal of Health and Physical Activity* 4: 67-71.
16. Budi S, Sri P, Wasilah R, dan Retno P (2011) Hubungan Indeks Massa Tubuh Dengan Densitas Mineral Tulang Pada Perempuan Dewasa Muda. *Pusat Penelitian dan Pengembangan Gizi dan Makanan, Badan Penelitian dan Pengembangan Kesehatan. Kemenkes RI* 34: 93-103.
17. Iqbal SI, Morch LS, Rosenzweig M, Dela F (2005) The Outcome of Bone Mineral Density Measurements on Patients Referred From General Practise. *Journal of Clinical Densitometry* 8: 178-182.

-
18. Herkulina F (2012) Hubungan Indeks Massa Tubuh dan Komposisi Tubuh dengan Densitas Massa Tulang Wanita Dewasa di Perwatusi Kota Pontianak. Electronic Tesis dan disertasi Universitas Gajah Mada.
 19. Kim SJ, Yang WG, Cho E, Park E (2012) Relationship between Weight, Body Mass Index and Bone Mineral Density of Lumbar Spine in Women. *J Bone Metab* 19: 95-102.
 20. Hubungan NA (2004) Tingkat Konsumsi Kalsium, Protein dan Status Gizi dengan Derajat Osteoporosis Pada lansia.
 21. Mendez JP, Mejia DR, Pedraza J, Varquez RMC, Soriano R, et al. (2013) Bone Mineral Density in Postmenopausal Mexican-Mestizo Women with Normal Body Mass Index, Overweight, or Obesity. *Menopause* 20: 568-72.
 22. Kalaselvi, Prabhu K, Ramesh M, Venkatesan V (2013) The Association of Serum Osteocalcin with the Bone Mineral Density in Post-Menopausal Women. *J Clin Diagn Res* 7: 814-816.

The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI), and Osteocalcin in Postmenopausal Women

ORIGINALITY REPORT

20%

SIMILARITY INDEX

10%

INTERNET SOURCES

19%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1

Øyen, Jannike, Clara Gram Gjesdal, Ottar Kjell Nygård, Stein Atle Lie, Haakon E. Meyer, Ellen Margrete Apalset, Per Magne Ueland, Eva Ringdal Pedersen, Øivind Midttun, Stein Emil Vollset, and Grethe S. Tell. "Smoking and Body Fat Mass in Relation to Bone Mineral Density and Hip Fracture: The Hordaland Health Study", PLoS ONE, 2014.

Publication

1%

2

Juan Pablo Méndez, David Rojano-Mejía, Ramón Mauricio Coral-Vázquez, Agustín Coronel et al. "Impact of genetic variants of IL-6, IL6R, LRP5, ESR1 and SP7 genes on bone mineral density in postmenopausal Mexican-Mestizo women with obesity", Gene, 2013

Publication

1%

3

Abdel-Aziz, M.M.. "Mutant p53 protein in the serum of patients with colorectal cancer: Correlation with the level of carcinoembryonic antigen and serum epidermal growth factor

1%

receptor", Cancer Detection and Prevention, 2009

Publication

4

"IUNS. 21st International Congress of Nutrition. Buenos Aires, Argentina, October 15-20, 2017: Abstracts", Annals of Nutrition and Metabolism, 2017

Publication

1 %

5

Lee, A. J., S. Hodges, and R. Eastell. "Measurement of Osteocalcin", Annals of Clinical Biochemistry An international journal of biochemistry and laboratory medicine

Publication

1 %

6

"ASBMR 22nd annual meeting", Journal of Bone and Mineral Research, 09/2000

Publication

1 %

7

Bone Densitometry in Clinical Practice, 2010.

Publication

1 %

8

"Poster presentations abstracts", Osteoporosis International, 2007

Publication

1 %

9

Kyong-Chol Kim, Dong-Hyuk Shin, Sei-Young Lee, Jee-Aee Im, Duk-Chul Lee. "Relation between Obesity and Bone Mineral Density and Vertebral Fractures in Korean Postmenopausal Women", Yonsei Medical Journal, 2010

Publication

1 %

10

Marina Cecelja, Michelle L. Frost, Tim D. Spector, Phil Chowienczyk. "Abdominal Aortic Calcification Detection Using Dual-Energy X-Ray Absorptiometry: Validation Study in Healthy Women Compared to Computed Tomography", *Calcified Tissue International*, 2013

Publication

1 %

11

Jaak Jürimäe, Priit Purge, Toivo Jürimäe, Serge P. von Duvillard. "Bone metabolism in elite male rowers: adaptation to volume-extended training", *European Journal of Applied Physiology*, 2006

Publication

1 %

12

Samuel D. Vasikaran. "Utility of Biochemical Markers of Bone Turnover and Bone Mineral Density in Management of Osteoporosis", *Critical Reviews in Clinical Laboratory Sciences*, 2008

Publication

1 %

13

N.A. Prastowo, S. Ali, Ignatio Rika Haryono. "A Population-based Study on Bone Mineral Density Using Dual-Energy X-Ray Absorptiometry (DEXA) in Postmenopausal Women in Jakarta, Indonesia", *International Journal of Osteoporosis and Metabolic Disorders*, 2018

Publication

1 %

14

M. Lateef. "Estimation of serum osteocalcin and telopeptide-C in postmenopausal osteoporotic females", Osteoporosis International, 07/14/2009

Publication

<1 %

15

idosi.org

Internet Source

<1 %

16

www.ncbi.nlm.nih.gov

Internet Source

<1 %

17

www.jorl.net

Internet Source

<1 %

18

Komang Agung Irianto, Moulli Edward, Angga Fiandana. "K-wire migration to unexpected site", International Journal of Surgery Open, 2018

Publication

<1 %

19

"New Innovative Technologies Changing Cytology. 19th International Congress of Cytology. Yokohama, Japan, May 29-June 1, 2016: Abstracts", Acta Cytologica, 2016

Publication

<1 %

20

"Abstracts of the 35th European Symposium on Calcified Tissues", Calcified Tissue International, 2008

Publication

<1 %

21

Phui-Ly Liew, Weu Wang, Yi-Chih Lee, Ming-Te Huang, Yang-Chu Lin, Wei-Jei Lee.

<1 %

"Gallbladder Disease among Obese Patients in Taiwan", Obesity Surgery, 2007

Publication

22

citeseerx.ist.psu.edu

Internet Source

<1 %

23

Osama Farouk, Dalia G. Mahran, Hatem G. Said, Mohamed M. Alaa, Amr Eisa, Hisham Imam, GZ Said. "Osteoporosis among hospitalized patients with proximal femoral fractures in Assiut University Trauma Unit, Egypt", Archives of Osteoporosis, 2017

Publication

<1 %

24

www.dovepress.com

Internet Source

<1 %

25

actaepidendocrinol.com

Internet Source

<1 %

26

Khuloud Bajbouj, Jasmin Shafarin, Hilda Allam, Mohamed Madkour et al. "Elevated Levels of Estrogen Suppress Hepcidin Synthesis and Enhance Serum Iron Availability in Premenopausal Women", Experimental and Clinical Endocrinology & Diabetes, 2018

Publication

<1 %

27

Dini Setiarsih, Bambang Wirjatmadi, Merryana Adriani. "Bone Density Status and Vitamin D and Calcium Concentrations in Pregnant and

<1 %

Non-Pregnant Women", Makara Journal of Health Research, 2016

Publication

-
- | | | |
|----|---|------|
| 28 | global.oup.com
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|--|------|
| 29 | Shun-zhi Liu, Li-fang Tian, Peng Xu, Gui-hua Zhuang, Fang Zheng, Juan Tian, Qi-Lan Ning, Bo-Feng Zhu, She-Min Lu, Hong Yan. "Analysis of correlation between blood biochemical indicators and bone mineral density of post-menopausal women", Molecular Biology Reports, 2010
Publication | <1 % |
|----|--|------|
-
- | | | |
|----|---|------|
| 30 | epublications.uef.fi
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 31 | www.ijmhs.net
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 32 | profdoc.um.ac.ir
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 33 | www.micromedex.com
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 34 | www.biomedcentral.com
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|--|------|
| 35 | "Poster Presentations - Clinical abstracts (ICOBR 2014)", IBMS BoneKEy, 2014.
Publication | <1 % |
|----|--|------|

36

"Free Communication Sessions 01-20 and Poster Sessions 01-16", International Dental Journal, 2015.

Publication

<1 %

37

Thijssen, Jos H. H.. "Gene polymorphisms involved in the regulation of bone quality", Gynecological Endocrinology, 2006.

Publication

<1 %

38

"ECTS Abstracts", Calcified Tissue International, 2003

Publication

<1 %

39

Shahzad, K.. "Analytic evaluation of the @b-human chorionic gonadotropin assay on the Abbott IMx and Elecsys2010 for its use in doping control", Clinical Biochemistry, 200711

Publication

<1 %

Exclude quotes

On

Exclude matches

Off

Exclude bibliography

On

The Correlation of Bone Mineral Density (BMD), Body Mass Index (BMI), and Osteocalcin in Postmenopausal Women

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5